

TEICHMÜLLER GEODESICS THAT DON'T HAVE A LIMIT IN $\mathcal{PMF}$

ANNA LENZHEN

ABSTRACT. We construct a Teichmüller geodesic which does not have a limit on the Thurston boundary of the Teichmüller space.

1. INTRODUCTION.

Let M be a closed surface of genus $g \geq 2$. Recall that the *Teichmüller space* $\mathcal{T}_g$ is the space of equivalence classes of conformal structures X on M , where $X_1 \sim X_2$ if there is a biholomorphic map from X_1 to X_2 which is isotopic to the identity on M . The *Teichmüller distance* between $X_1, X_2 \in \mathcal{T}_g$ is defined by $d(X_1, X_2) = \frac{1}{2} \log K$, where K is the smallest number such that there is a homeomorphism homotopic to the identity on M which is a K -quasiconformal map between X_1 and X_2 . A *holomorphic quadratic differential* q on a Riemann surface X assigns to each chart (U_α, z_α) of X a holomorphic function q_{z_α} with the property $q_{z_\beta}(z_\beta) \left(\frac{dz_\alpha}{dz_\beta} \right)^2 = q_{z_\alpha}(z_\alpha)$ in $U_{z_\alpha} \cap U_{z_\beta}$. It determines a flat metric with the length element $|q(z)|^{1/2} |dz|$. The vertical trajectories of q are the arcs along which $q(z) dz^2 < 0$, and the horizontal trajectories are the arcs where $q(z) dz^2 > 0$. Hence every quadratic differential q determines a pair of transverse measured foliations: the *vertical* foliation, where the leaves are the vertical trajectories, and the *horizontal* foliation, with leaves being the horizontal trajectories of q . The norm or area of q is defined by $\|q\| = \int_M |q(z)| |dz|^2$. In a neighborhood of every regular point P of q one can introduce a local parameter w , in terms of which q is identically equal to 1. This parameter, called the *natural parameter* of q near P , is determined by the integral $w = \int_X \sqrt{q(z)} dz$, uniquely up to a transformation $w \rightarrow \pm w + \text{const}$. The space Q_0 of all holomorphic quadratic differentials on X is a $6g - 6$ dimensional vector space.

Each direction at a point X in $\mathcal{T}_g$ is associated to a quadratic differential q on X . The Teichmüller geodesic determined by q consists of the one parameter family $\{X_t\}$ where X_t is the conformal structure X is sent to by a map which can be described locally by $x \rightarrow e^{t/2}x$, $y \rightarrow e^{-t/2}y$ where $w = x + iy$ is the natural parameter, so that the transverse measure on the vertical (horizontal) foliation of q is scaled by $e^{t/2}$ ($e^{-t/2}$).

Let $\mathcal{S}$ be the set of homotopy classes of simple closed curves on M with the discrete topology; R_+^S is given the product topology and PR_+^S is the corresponding projective space. The Teichmüller space $\mathcal{T}_g$ can be identified with the space of equivalence classes of hyperbolic metrics ρ on M of constant curvature -1 , where $\rho_1 \sim \rho_2$ if there exists an isometry from (M, ρ_1) to (M, ρ_2) isotopic to the identity. The map $\mathcal{T}_g \mapsto PR_+^S$ defined by $\rho \mapsto (\gamma \mapsto \ell_\rho(\gamma))$, where $\ell_\rho(\gamma)$ is the length of the unique geodesic in the hyperbolic metric

in the class of γ , is injective. It is called the Thurston embedding of Teichmüller space. The boundary of $\mathcal{T}_g$ in PR_+^S is the sphere $\mathcal{PMF}$ of projective measured foliations on M . The union of $\mathcal{T}_g$ and $\mathcal{PMF}$ is denoted by $\hat{\mathcal{T}}_g$ and is called Thurston compactification. It is homeomorphic to a closed $6g - 6$ dimensional ball, where $\mathcal{PMF}$ is the $6g - 7$ dimensional sphere.

Masur (see [5]) proved that in almost every direction through every point geodesic rays converge in $\hat{\mathcal{T}}_g$. In particular, if the vertical foliation F of a quadratic differential q is uniquely ergodic, then the geodesic ray corresponding to q converges to the class of F in $\mathcal{PMF}$. In the same paper he showed that, if all leaves of F are closed, then the ray also converges. In this paper we show that not every Teichmüller geodesic ray converges to a single measured foliation. More precisely, we construct a geodesic ray in $\mathcal{T}_2$ which does not have a limit in $\mathcal{PMF}$. The quadratic differential that determines the direction in our construction has non-minimal vertical foliation. The question of convergence in the general case, in particular when the vertical foliation is minimal but not uniquely ergodic, remains open.

Construction. Start with a square torus, i.e. the unit square with lower left vertex at $(0, 0)$, with opposite sides identified. Cut the torus along a line segment (call it a slit) of length $0 < s < 1$ and a slope $\theta_1 > 0$. Take a second copy of the torus, cut it along a slit of same length s and a slope $\theta_2 > 0$. Rotate the squares counterclockwise so that the slits are vertical. Now identify the positive side of the slit on one copy with the negative side on the other copy. This results in a quadratic differential q_{θ_1, θ_2} with 2 zeroes of order 2 on a surface X of genus 2. X is partitioned into 2 sheets S_1 and S_2 separated from each other by the union of the two slits. If both θ_1 and θ_2 are irrational, then the vertical foliation of q_{θ_1, θ_2} has one closed leaf, which is the union of the two slits, and all other leaves are dense in S_1 or S_2 . Let $\{X_t\}$ be the Teichmüller geodesic ray from X determined by the differential q_{θ_1, θ_2} .

Recall that each $x \in \mathbb{R}$ admits a continued fraction expansion of the form $x = a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \dots}}$, with $a_0 \in \mathbb{Z}$, $a_i \in \mathbb{N}$, $i \geq 1$ and that $x \in \mathbb{R} - \mathbb{Q}$ iff infinitely many a_i 's are

nonzero. We will also use the notation $x = [a_0; a_1, a_2, \dots]$. We will call a_i 's the *elements* of x .

Our main result is

Theorem 1. *Suppose $\theta_1 \in \mathbb{R} - \mathbb{Q}$ has bounded elements $a_{1,n} \geq 3$, $n \geq 1$, and $\theta_2 \in \mathbb{R} - \mathbb{Q}$ has unbounded elements $a_{2,n} \geq 3$, $n \geq 1$. Let $\{X_t\}$ be the Teichmüller geodesic ray constructed as above. Then $\{X_t\}$ does not converge in $\hat{\mathcal{T}}_2$.*

Idea of the proof. By the definition of Thurston compactification, a sequence $\{X_t\}$ in Teichmüller space converges to projective measured foliation F in $\mathcal{PMF}$ if, for any $\alpha_1, \alpha_2 \in \mathcal{S}$ we have $\frac{\ell_t(\alpha_1)}{\ell_t(\alpha_2)} \xrightarrow{t \rightarrow \infty} \frac{i(\alpha_1, F)}{i(\alpha_2, F)}$. Here $\ell_t(\alpha_i)$ is the length of the shortest curve in the hyperbolic metric of X_t which is homotopic to a curve α_i , and $i(\alpha_i, F)$ denotes the measure of α_i with

respect to F . Hence, to prove that the geodesic does not have a limit in $\mathcal{PMF}$, it suffices to find a pair of simple closed curves α_1 and α_2 on M and show that the limit $\lim_{t \rightarrow \infty} \frac{\ell_t(\alpha_1)}{\ell_t(\alpha_2)}$ does not exist. In our proof α_i is the curve represented by the vector $(1, 0)$ on S_i , $i = 1, 2$.

To prove the theorem, we need to be able to estimate hyperbolic lengths of simple closed curves on X_t for large values of t . The geodesic ray $\{X_t\}$ is constructed using the flat metric. However, in general there is no easy way to get a good estimate of the hyperbolic length of a curve. Our method is to identify the shortest and the second shortest curves in the flat metric on each torus (we think of the surface X_t as a surface glued out of two tori S_1 and S_2), and estimate their hyperbolic length. This information, together with the notion of intersection number, makes it possible to find good lower and upper bounds on the lengths of the curves α_1 and α_2 .

The idea behind the choice of θ_1 and θ_2 is as follows. We will show (Lemma 1) that at any time the shortest curve on S_i is a curve whose slope is a convergent of θ_i . Moreover, we will prove (Lemma 3) that there is a sequence of times t_k and a sequence of elements $a_{2,n(t_k)} \rightarrow \infty$ of θ_2 for which the length (flat, extremal and hyperbolic) of the shortest curve on S_2 goes to 0, while on the other side the length of the shortest curve stays bounded below. By the Collar Lemma (see [2]), each time one crosses an extremely short curve, one has to cross a collar of width approximately $\log \frac{1}{\ell(\text{short curve})}$. Hence one would think that the curve α_2 must become exceedingly long compared to the curve α_1 . However, we will see that the curve α_1 intersects a curve of bounded length on S_1 significantly more than the curve α_2 crosses the long collar in S_2 . It turns out that the ratio of the lengths $\ell_{t_k}(\alpha_1)$ and $\ell_{t_k}(\alpha_2)$ is at least $\frac{\sqrt{a_{2,n(t_k)}}}{\log a_{2,n(t_k)}}$ and is unbounded. On the other hand, when there are no short curves, the ratio of the lengths of α_1 and α_2 stays bounded, and therefore we conclude that the limit $\lim_{t \rightarrow \infty} \frac{\ell_t(\alpha_1)}{\ell_t(\alpha_2)}$ does not exist.

The main result is proven in Section 4. In Section 2 we consider the shortest curves in the flat metric. Estimates of the hyperbolic lengths of these curves are made in Section 3.

Acknowledgements. The author would like to thank her thesis advisor Howard Masur for his excellent guidance and support.

2. FLAT METRIC

In this section we focus on one torus only, ignoring the slit and the other torus. Let $\theta = [a_0; a_1, a_2, a_3, \dots]$ be any positive irrational number, with $a_i \geq 3$. The k -th convergent of θ is the reduced fraction $\frac{p_k}{q_k} = a_0 + \frac{1}{a_1 + \frac{1}{\dots + \frac{1}{a_k}}}$. A few standard facts (see for

example [4]) about continued fractions are:

$$(1a) \quad p_{n+1} = a_{n+1}p_n + p_{n-1} \text{ and } q_{n+1} = a_{n+1}q_n + q_{n-1}$$

$$(1b) \quad \frac{1}{q_n + q_{n+1}} \leq |p_n - \theta q_n| \leq \frac{1}{q_{n+1}}$$

$$(1c) \quad \frac{p_{2n}}{q_{2n}} \nearrow \theta \text{ and } \frac{p_{2n+1}}{q_{2n+1}} \searrow \theta$$

$$(1d) \quad |p_{n+1}q_n - q_{n+1}p_n| = 1$$

Consider a standard lattice $\mathbb{Z}^2$ in $\mathbb{R}^2$. Let g_t^θ be a map which at time $t \geq 0$ is a rotation by an angle of $\frac{\pi}{2} - \tan^{-1} \theta$, followed by a horizontal stretch by a factor of $e^{t/2}$ and vertical contraction by $e^{t/2}$. For every t we get a new lattice in $\mathbb{R}^2$. We will refer to the image of any vector $(q, p) \in \mathbb{Z}^2$ under the map g_t^θ as (q, p) -vector or (q, p) -curve at time t .

Notation. To simplify our presentation we use $\approx$, Θ and O defined as follows: for two sequences $x_n > 0, y_n > 0$, $x_n \approx y_n$ means $\frac{x_n}{y_n} \xrightarrow{n \rightarrow \infty} 1$; $x_n = O(y_n)$ iff $\sup \frac{x_n}{y_n} < \infty$; $x_n = \Theta(y_n)$ iff $x_n = O(y_n)$ and $y_n = O(x_n)$.

Lemma 1. Suppose a (q, p) -vector has the shortest euclidean length at some time t . Then $\frac{p}{q}$ is a convergent for θ , i.e. $p = p_n$ and $q = q_n$ for some n . Also at time $T_n = \log \frac{p_n \theta + q_n}{|q_n \theta - p_n|}$, the (q_n, p_n) -vector is the shortest. For $t \in [T_n, T_{n+1}]$ the shortest vector is either (q_n, p_n) or (q_{n+1}, p_{n+1}) .

Proof. We can assume that

$$\frac{p_2}{q_2} < \frac{p}{q} < \theta < \frac{p_1}{q_1}$$

If $\frac{p}{q}$ is not a convergent of θ , then there is a unique n such that $\frac{p_n}{q_n} < \frac{p}{q} < \frac{p_{n+2}}{q_{n+2}}$. We claim that the (q_{n+1}, p_{n+1}) -vector is always shorter than the vector (q, p) . At time t the image of (q, p) is $g_t^\theta(q, p) = \frac{q + \theta p}{e^{t/2} \sqrt{1 + \theta^2}}(0, 1) + \frac{e^{t/2}(q\theta - p)}{\sqrt{1 + \theta^2}}(1, 0)$. The euclidean length of $g_t^\theta(q, p)$ satisfies

$$l((q, p)) = \frac{1}{1 + \theta^2} \left(\frac{(q + \theta p)^2}{e^t} + e^t(p - \theta q)^2 \right).$$

Since $\frac{p_n}{q_n} < \frac{p}{q} < \frac{p_{n+1}}{q_{n+1}}$ and $p_{n+1}q_n - q_{n+1}p_n = 1$, it is easy to see that $p \geq p_n + p_{n+1}$ and $q \geq q_n + q_{n+1}$. Therefore

$$(2) \quad q + \theta p > q_{n+1} + \theta p_{n+1}$$

Also

$$(3) \quad |p - \theta q| = q \left| \frac{p}{q} - \theta \right| > q \left| \frac{p}{q} - \frac{p_{n+2}}{q_{n+2}} \right| > \frac{1}{q_{n+2}} > |p_{n+1} - \theta q_{n+1}|$$

The claim now follows from (2) and (3) and we conclude that $\frac{p}{q}$ is a convergent of θ .

The function $l((q_n, p_n)_t)$ reaches its minimum of $2 \frac{(q_n + p_n \theta) |q_n \theta - p_n|}{1 + \theta^2}$ at $T_n = \log \frac{p_n \theta + q_n}{|q_n \theta - p_n|}$. We

have for all $k < n$

$$l_{T_n}((q_k, p_k))^2 > \frac{(q_k\theta - p_k)^2 e^{T_n}}{1 + \theta^2} = \frac{(q_k\theta - p_k)^2 (p_n\theta + q_n)}{(1 + \theta^2)|q_n\theta - p_n|}$$

and for all $k > n$

$$l_{T_n}((q_k, p_k))^2 > \frac{(q_k + p_k\theta)^2}{(1 + \theta^2)e^{T_n}} = \frac{(q_k + p_k\theta)^2 |q_n\theta - p_n|}{(1 + \theta^2)(p_n\theta + q_n)}$$

Then

$$l_{T_n}^2((q_k, p_k)) > \frac{q_{n+1}}{4q_n}, \quad k < n$$

and

$$l_{T_n}^2((q_k, p_k)) > \frac{q_{n+1}}{2q_n}, \quad k > n$$

On the other hand

$$2\frac{q_n}{q_n + q_{n+1}} \leq l_{T_n}^2((q_n, p_n)) \leq 2\frac{q_n}{q_{n+1}}$$

The last statement follows from (1b) and the fact that $p_n\theta + q_n \approx q_n(1 + \theta^2)$. We assumed that the elements of θ satisfy $a_k > 2$, so we have $l_{T_n}((q_n, p_n)) < \min_{k \neq n} \{l_{T_n}((q_k, p_k))\}$.

To verify the last claim of the lemma, we notice that, perhaps except for the first few




FIGURE 1

terms, the sequence $\{T_n\}$ is strictly increasing. Also we see that, for each n , the function $l_t^2((q_n, p_n))$ is strictly convex and that the graphs of $l_t^2((q_n, p_n))$ and $l_t^2((q_k, p_k))$ intersect once. Since $n < k$ implies $l_0^2((q_n, p_n)) < l_0^2((q_k, p_k))$, we see that at time $t \in [T_n, T_{n+1}]$ only (q_n, p_n) or (q_{n+1}, p_{n+1}) can be the shortest.

□

To find the second shortest vector is a more complicated task. There are more curves which can become second shortest then just the ones whose slopes are convergents of θ .

Lemma 2. *Suppose that a (q, p) -curve is second shortest at time t , and (q_n, p_n) is the shortest. Then either $\frac{p}{q}$ is a convergent of θ (namely, (q_{n-1}, p_{n-1}) or (q_{n+1}, p_{n+1})) or the following holds*

$$q_{n-1} + q_n \leq q < q_{n+1}$$

and

$$p_{n-1} + p_n \leq p < p_{n+1}$$

Proof. Suppose $\frac{p}{q}$ is not a convergent of θ . Assume that $\frac{p}{q} < \theta$. Then there is a (unique) k such that $\frac{p_{k-1}}{q_{k-1}} < \frac{p}{q} < \frac{p_{k+1}}{q_{k+1}}$. Using the argument from Lemma 1 we can show that the (q_k, p_k) -curve is shorter than (q, p) . Therefore $k = n$. Since the shortest and the second shortest curves intersect once, $p_n q - p q_n = 1$. Also $p_n q_{n-1} - q_n p_{n-1} = 1$. Putting these two things together we get

$$p_n(q - q_{n-1}) - q_n(p - p_{n-1}) = 0$$

This implies that $\frac{p_n}{q_n} = \frac{p - p_{n-1}}{q - q_{n-1}}$. Hence $p = ap_n + p_{n-1}$ and $q = aq_n + q_{n-1}$, where $a \in \mathbb{Q}$. It is easy to see that $a < a_{n+1}$: if $a \geq a_{n+1}$ then

$$\frac{p}{q} - \frac{p_{n+1}}{q_{n+1}} = \frac{ap_n + p_{n-1}}{aq_n + q_{n-1}} - \frac{a_{n+1}p_n + p_{n-1}}{a_{n+1}q_n + q_{n-1}} = \frac{(a_{n+1} - a)(p_{n-1}q_n - q_{n-1}p_n)}{(aq_n + q_{n-1})(a_{n+1}q_n + q_{n-1})} \geq 0$$

which is impossible. Therefore $p < p_{n+1}$ and $q < q_{n+1}$. Also $\frac{p_{n-1}}{q_{n-1}} < \frac{p}{q} < \frac{p_n}{q_n}$ and $p_n q_{n-1} - q_n p_{n-1} = 1$ imply $p \geq p_{n-1} + p_n$ and $q \geq q_{n-1} + q_n$. A similar argument works if $\frac{p}{q} > \theta$.

If $\frac{p}{q}$ is a convergent of θ , then Lemma 1 implies that (q, p) is either (q_{n-1}, p_{n-1}) or (q_{n+1}, p_{n+1}) . $\square$

3. HYPERBOLIC METRIC.

In the previous section we considered shortest vectors on a torus. It follows from Lemma 1 that the vectors on S_i with smallest euclidean length are those whose slopes are n -th convergents of θ_i , which we denote by $\frac{p_{i,n}}{q_{i,n}}$. We claim that the same result holds for the flat surface glued from the two tori S_1 and S_2 , each with a slit $(0, s)$. The point is that, for each n , the curve $(q_{i,n}, p_{i,n})$ on S_i can be made disjoint from the slit.

Hence at any time we know the short curves in S_1 and S_2 . We want to estimate the hyperbolic length of these curves. To do so, we first prove a claim about extremal lengths of curves in S_i which get short. We will denote by $Ext_t(\alpha)$ the extremal length at time t of a family of curves homotopic to a curve α .

Lemma 3. *Let α be a $(q_{i,n}, p_{i,n})$ -curve on S_i , $i = 1, 2$. Then the extremal length of α at time t satisfies*

$$\frac{Ext_t(\alpha)}{l_t^2(\alpha)} \xrightarrow{n, t \rightarrow \infty} 1$$

Proof. By definition of extremal length $Ext(\alpha) \geq \frac{\inf_{\beta \sim \alpha} (\int_{\beta} \rho |dz|)^2}{A(\rho)}$ for any metric of the form $\rho |dz|$ where $\rho \geq 0$ is Borel measurable. Assume α is a $(p_{1,n}, q_{1,n})$ -curve on S_1 . Let $\rho_t |dz|$ be




FIGURE 2. $\rho_t |dz|$ coincides with the flat metric in the shaded area

the metric which coincides with the flat metric on S_1 , and on the set of points in S_2 which are at most $se^{-t/2}$ (the length of the slit) away from the slit (see figure 2). We need to find the shortest curve with respect to this metric in the homotopy class of α . We claim that a geodesic with respect to $\rho_t |dz|$ is contained in S_1 , i.e. it does not cross the slit. The point is that any curve β homotopic to α which crosses the slit will contain an arc in S_2 with endpoints on the slit. The arc is longer than the line segment connecting the endpoints. Hence there is a curve in the class of α which is shorter than β . Since α is a geodesic in the flat metric, the length of the shortest curve homotopic to α with respect to $\rho_t |dz|$ is at least the flat length of α , which is $\sqrt{\frac{1}{1+\theta_1^2} \left(\frac{(q_{1,n} + \theta_1 p_{1,n})^2}{e^t} + e^t (p_{1,n} - \theta_1 q_{1,n})^2 \right)}$. We then have




FIGURE 3

$$(4) \quad Ext_t(\alpha) \geq \frac{1}{1 + \theta_1^2} \frac{\frac{(q_{1,n} + \theta_1 p_{1,n})^2}{e^t} + e^t (p_{1,n} - \theta_1 q_{1,n})^2}{1 + (\pi + 2)s^2 e^{-t}}$$

On the other hand, the extremal length of a simple closed curve α can be defined as $\sup_C \frac{1}{Mod(C)}$ where the supremum is taken over all cylinders C with α a core curve, and where $Mod(C)$ denotes the modulus of a cylinder C .

So we need to find a cylinder for a good upper bound. We consider a cylinder swept out by curves parallel to α which avoid the slit. Curves parallel to α and crossing the slit make up a parallelogram spanned by the vectors representing the slit and α . At time t the slit is represented by a vector $u = (0, \frac{s}{e^{t/2}})$, and α is represented by $v = \frac{q_{1,n} + \theta p_{1,n}}{e^{t/2} \sqrt{1 + \theta_1^2}}(0, 1) + \frac{e^{t/2} |q_{1,n} \theta - p_{1,n}|}{\sqrt{1 + \theta_1^2}}(1, 0)$. Then the area of our cylinder is

$$1 - |u \times v| = 1 - \frac{s}{e^{t/2}} \frac{e^{t/2} |q_{1,n} \theta - p_{1,n}|}{\sqrt{1 + \theta_1^2}} = 1 - \frac{s |q_{1,n} \theta - p_{1,n}|}{\sqrt{1 + \theta_1^2}}$$

The length of the cylinder is $\|v\|$, the height is

$$\frac{area}{length} = \frac{1 - \frac{s |q_{1,n} \theta - p_{1,n}|}{\sqrt{1 + \theta_1^2}}}{\|v\|}$$

Thus the modulus is

$$\frac{1 - \frac{s |q_{1,n} \theta - p_{1,n}|}{\sqrt{1 + \theta_1^2}}}{\|v\|^2}$$

It follows that

$$(5) \quad Ext_t(\alpha) \leq \frac{1}{1 + \theta_1^2} \left(\frac{(q_{1,n} + \theta_1 p_{1,n})^2}{e^t} + e^t (p_{1,n} - \theta_1 q_{1,n})^2 \right) \frac{1}{1 - \frac{s |q_{1,n} \theta - p_{1,n}|}{\sqrt{1 + \theta_1^2}}}$$

The claim of the lemma now follows from (5), (4) and (1b). $\square$

Corollary 1. *Suppose θ_i is such that $a_{i,n} \rightarrow \infty$ for $i = 1$ or $i = 2$. Then at time $T_n = \log \frac{p_{i,n} \theta_i + q_{i,n}}{|q_{i,n} \theta_i - p_{i,n}|}$ the hyperbolic length of a curve $\alpha = (q_{i,n}, p_{i,n})$ satisfies $\ell_{T_n}(\alpha) = \Theta(\frac{1}{a_{i,n+1}})$.*

Proof. By Lemma 3 we have $Ext_{T_n}(\alpha) \approx l_{T_n}^2(\alpha)$. Furthermore, the assumption $a_{i,n} \rightarrow \infty$ and (1b) imply $l_{T_n}^2(\alpha) \approx \frac{2q_{i,n}}{q_{i,n+1}} \approx \frac{2}{a_{i,n+1}}$. On the other hand, by Proposition 1 and Corollary 3 in [6] that $2e^{-\ell(\gamma)/2} \leq \frac{\ell(\gamma)}{Ext(\gamma)} \leq \pi$, where $Ext(\gamma)$ is the extremal length and $\ell(\gamma)$ - the hyperbolic length of any simple closed curve γ . In particular, if the extremal length of α becomes very small, then so does the hyperbolic length, and their ratio is bounded above and below. Therefore, the hyperbolic length of α at time T_n satisfies $\ell_{T_n}(\alpha) = \Theta(\frac{1}{a_{i,n+1}})$. $\square$

4. PROOF OF THE MAIN THEOREM.

We begin by choosing a subsequence $a_{2,n_k} \rightarrow \infty$ such that $a_{2,n_k} \rightarrow \infty$.
Let for $k \in \mathbb{Z}, k \geq 1$

$$t_{2k} = \log \frac{p_{2,n_k-1}\theta_2 + q_{2,n_k-1}}{|q_{2,n_k-1}\theta_2 - p_{2,n_k-1}|}$$

$$t_{2k+1} = \log((1 + \theta_2^2)(q_{2,n_k})^2)$$

We'll show that $\lim_{i \rightarrow \infty} \frac{\ell_{t_i}(\alpha_1)}{\ell_{t_i}(\alpha_2)}$ does not exist. By $\alpha_{1,1}(i)$ and $\alpha_{1,2}(i)$ (respectively $\alpha_{2,1}(i)$ and $\alpha_{2,2}(i)$) we denote curves on S_1 (respectively S_2) which are the first and second shortest in the flat metric at time t_i .

i = 2k. We have for sufficiently large k

$$\ell_{t_{2k}}(\alpha_2) \leq \ell_{t_{2k}}(\alpha_{2,1}(2k))i(\alpha_{2,2}(2k), \alpha_2) + \ell_{t_{2k}}(\alpha_{2,2}(2k))i(\alpha_{2,1}(2k), \alpha_2)$$

Lemma 1 implies that $\alpha_{2,1}(2k)$ is a $(q_{2,n_k-1}, p_{2,n_k-1})$ -curve. Thus $i(\alpha_{2,1}(2k), \alpha_2) = p_{2,n_k-1}$. Corollary 1 implies that $\ell_{t_{2k}}(\alpha_{2,1}(2k)) = \Theta(\frac{1}{a_{2,n_k}})$. If the second shortest curve is a (q, p) -curve, then by Lemma 2 we have $p \leq p_{2,n_k}$. Therefore $i(\alpha_{2,2}(2k), \alpha_2) = p \leq p_{2,n_k}$.

To estimate $\ell_{t_{2k}}(\alpha_{2,2}(2k))$, consider the flat cylinder from Lemma 3 with $\alpha_{2,1}(2k)$ the core curve. Its area (with respect to the flat metric) is $1 - s|q_{2,n_k-1}\theta_2 - p_{2,n_k-1}|$ and the modulus is $m = \frac{1 - s|q_{2,n_k-1}\theta_2 - p_{2,n_k-1}|}{l_{t_{2k}}^2(\alpha_{2,1}(2k))}$. It is conformally equivalent to the annulus $A = \{z \in \mathbb{C} | e^{-2\pi m} < |z| < 1\}$ with the hyperbolic metric

$$\rho_k(z) = \frac{-\pi|dz|}{|z|(2m) \sin(\log|z|/(2m))}$$

Since $\alpha_{2,2}(2k)$ is second shortest in the flat metric, it crosses any radius of the annulus at most once (otherwise there is a shorter curve in flat metric). From Lemma 3 in [5] we know that given $\epsilon > 0$ there exists $0 < \delta < 1$ such that for k sufficiently large the hyperbolic metric $\rho_{t_{2k}}(z)$ on $X_{t_{2k}}$ and the metric $\rho_k(x)$ satisfy

$$(6) \quad \left| \frac{\rho_{t_{2k}}(z)}{\rho_k(z)} - 1 \right| < \epsilon, \quad z \in A_\delta = \{e^{-2\pi m}/\delta < |z| < \delta\}$$

Hence for large enough k the hyperbolic length ℓ_1 of the part of $\alpha_{2,2}(2k)$ inside A_δ is approximately the length of the radius of A_δ with respect to ρ_k , more precisely

$$\ell_1 = \left(\log \frac{1 + \cos \frac{\log 1/\delta}{2m}}{1 - \cos \frac{\log 1/\delta}{2m}} \right) (1 + \epsilon) + C = \Theta\left(\log \frac{2m}{\log 1/\delta}\right) = \Theta(\log a_{2,n_k})$$

We claim that outside the annulus A_δ the hyperbolic length ℓ_2 of $\alpha_{2,2}(2k)$ is bounded above. Consider the family Γ of arcs in the complement of A_δ connecting the boundary components of the annulus. Clearly the subarc of $\alpha_{2,2}(2k)$ belongs to Γ . We claim that the extremal length of Γ is bounded. This follows from the fact that Γ contains a subset




FIGURE 4

P of arcs which connect opposite sides of a parallelogram of height

$$h = \frac{s|q_{2,n_k-1}\theta_2 - p_{2,n_k-1}|}{l_{t_{2k}}(\alpha_{2,1}(2k))} + \frac{1}{\pi} l_{t_{2k}}(\alpha_{2,1}(2k)) \log \frac{1}{\delta} = \Theta(l_{t_{2k}}(\alpha_{2,1}(2k)))$$

and length at least

$$l = l_{t_{2k}}(\alpha_{2,1}(2k)) - se^{-t_{2k}/2} = \Theta(l_{t_{2k}}(\alpha_{2,1}(2k)))$$

Hence the extremal length $Ext(P) = O(1)$. But $P \subset \Gamma$ implies $Ext(P) \geq Ext(\Gamma)$. Therefore the shortest in hyperbolic metric element of Γ is bounded. Since ℓ_2 differs from it by an additive constant, it is also bounded.

We then have

$$(7) \quad \ell_{t_{2k}}(\alpha_2) = O(\log(a'_{n_k})p'_{n_k-1})$$

Now we want to estimate $\ell_{t_{2k}}(\alpha_1)$. Recall that θ_1 is such that the elements satisfy $\sup_i a_{1,i} < \infty$. Then Lemma 1 and Lemma 3 imply that $Ext_{t_{2k}} \alpha_{1,1}(2k) = \Theta(1)$. Hence $\ell_{t_{2k}}(\alpha_{1,1}(2k)) = \Theta(1)$. Every time α_1 crosses $\alpha_{1,1}(2k)$, it has to cross the collar around $\alpha_{1,1}(2k)$, which by the Collar Lemma (see [2]) has length of at least $2\text{arcsinh}\left(\frac{1}{\sinh(\ell_{t_{2k}}(\alpha_{1,1}(2k))/2)}\right) = \Theta(1)$. Hence

$$(8) \quad \ell_{t_{2k}}(\alpha_1) \geq \Theta(i(\alpha_{1,1}(2k), \alpha_1))$$

If $\alpha_{1,1}(2k)$ is a (q_{1,j_k}, p_{1,j_k}) -curve, then $i(\alpha_{1,1}(2k), \alpha_1) = p_{1,j_k} \approx q_{1,j_k}\theta_1$. By definition $\alpha_{1,1}(2k)$ is the shortest curve on S_1 at time t_{2k} , therefore it follows from Lemma 1 that the curve $(q_{1,j_k+1}, p_{1,j_k+1})$ has not reached its minimal length yet, i.e.

$$t_{2k} \leq \log \frac{p_{1,j_k+1}\theta_1 + q_{1,j_k+1}}{|q_{1,j_k+1}\theta_1 - p_{1,j_k+1}|}$$

Hence for sufficiently large k , using (1b), we get

$$q_{2,n_k}q_{2,n_k-1}(1+\theta_2^2) \leq \frac{p_{2,n_k-1}\theta + q_{2,n_k-1}}{|q_{2,n_k-1}\theta - p_{2,n_k-1}|} \leq \frac{p_{1,j_k+1}\theta_1 + q_{1,j_k+1}}{|q_{1,j_k+1}\theta_1 - p_{1,j_k+1}|} \leq (1+\theta_1^2)q_{1,j_k+1}(q_{1,j_k+1} + q_{1,j_k+2})$$

We then have $q_{1,j_k} = \Theta(q_{2,n_k-1}\sqrt{a_{2,n_k}})$. It follows that

$$(9) \quad \ell_{t_{2k}}(\alpha_1) \geq \Theta(q_{2,n_k-1}\sqrt{a_{2,n_k}})$$

Putting together (7) and (9) we get

$$(10) \quad \frac{\ell_{t_{2k}}(\alpha_1)}{\ell_{t_{2k}}(\alpha_2)} \geq \frac{\Theta(q_{2,n_k-1}\sqrt{a_{2,n_k}})}{\Theta(\log(a_{2,n_k})p_{2,n_k-1})} \xrightarrow{n \rightarrow \infty} \infty$$

$\mathbf{i} = 2\mathbf{k} + 1$. We will show that $\sup_n \frac{\ell_{t_{2k+1}}(\alpha_1)}{\ell_{t_{2k+1}}(\alpha_2)}$ is bounded. Recall that $t_{2k+1} = \log((1 + \theta_2^2)(q_{2,n_k})^2)$.

It is easy to see that $t_{2k} < t_{2k+1} < t_{2k+2}$. Hence by Lemma 1 $\alpha_{2,1}(2k)$ is either $(q_{2,n_k-1}, p_{2,n_k-1})$ - or (q_{2,n_k}, p_{2,n_k}) - curve. By Lemma 3 both $(q_{2,n_k-1}, p_{2,n_k-1})$ and (q_{2,n_k}, p_{2,n_k}) have at time t_{2k+1} extremal (and hyperbolic) length bounded above and below. Hence

$$(11) \quad \ell_{t_{2k+1}}(\alpha_2) \geq \Theta(p_{2,n_k})$$

On the other hand $\alpha_1(2k+1)$ and $\alpha_2(2k+1)$ also have bounded extremal length, therefore

$$\ell_{t_{2k+1}}(\alpha_1) = O(i(\alpha_{1,1}(2k+1), \alpha_1)) + O(i(\alpha_{1,2}(2k+1), \alpha_1))$$

To estimate $i(\alpha_{1,1}(2k+1), \alpha_1)$ we argue similarly to the case when $i = 2k$. If $\alpha_{1,1}(2k+1)$ is a (q_{1,j_k}, p_{1,j_k}) - curve, then Lemma 1 implies

$$t_{2k+1} \geq \log \frac{p_{1,j_k-1}\theta_1 + q_{1,j_k-1}}{|q_{1,j_k-1}\theta_1 - p_{j_k-1}|}$$

Hence, using (1b), for sufficiently large k we obtain

$$(1 + \theta_2^2)q_{1,n_k}^2 \geq \frac{p_{1,j_k-1}\theta_1 + q_{1,j_k-1}}{|q_{1,j_k-1}\theta_1 - p_{j_k-1}|} \geq q_{1,j_k-1}(1 + \theta_1^2)q_{1,j_k}$$

Thus $i(\alpha_{1,1}(2k+1), \alpha_1) = p_{1,j_k} = \Theta(q_{1,j_k}) = O(q_{2,n_k})$. Lemma 2 implies that

$$i(\alpha_{1,2}(2k+1), \alpha_1) \leq p_{1,j_k+1} = \Theta(p_{1,j_k})$$

We then have

$$(12) \quad \ell_{t_{2k+1}}(\alpha_1) = O(q_{2,n_k})$$

Putting (11) and (12) together we obtain

$$(13) \quad \sup_n \frac{\ell_{t_{2k+1}}(\alpha_1)}{\ell_{t_{2k+1}}(\alpha_2)} < \infty$$

REFERENCES

- [1] W Abikoff *Degenerating families of Riemann surfaces* Ann. of Math. (2) 105 (1977), no. 1, 29–44.
- [2] P Buser *Geometry and spektra of compact Riemann surfaces*, Progress in mathematics; v. 106, Birkhauser, Boston, 1992
- [3] Y Iwayoshi and M Taniguchi *An Introduction to Teichmüller Theory*, Springer-Verlag, 1992. MR 94b:32031
- [4] A Khinchin *Continued fractions*, Dover Publications, INC. Mineola, New York, 1997
- [5] H Masur *Two boundaries of Teichmüller space*, Duke Math., J 49, no. 1 (1982), 183-190
- [6] B Maskit *Comparison of hyperbolic and extremal lengths*, Ann. Acad. Sci. Fenn. Ser. A I Math. 10 (1985), 381-386. MR 87c:30062

- [7] G McShane, I Rivin *A norm on homology of surfaces and counting simple geodesics* eprint arXiv:math/0005222 (May 2000)